

1. Dressing Care:

Change your dressing two days after surgery. Some light bleeding is normal. Showering is permitted 2 days after surgery & the incision may be washed with soap & water, but no submerging or bathing is allowed until 2 weeks. Keep a clean dressing on the incision until your follow-up appointment.

2. Medication:

You will be given a painkiller _____ to be taken 1-2 pills every 4-6 hours as needed. Do not drink or drive while on pain medication. Do not take additional Tylenol based pain medications while on this medication.

3. Activity:

You will be placed in a sling after surgery. Only gentle passive pendulum range of motion is allowed in the sling with no external rotation or abduction of the arm. No other motion or activity is allowed at this time as this could result in tearing the repair.

Therapy will be started at 4-6 weeks to work on restoring motion and strengthening of the shoulder, progressing to full activity as tolerated. Expected time to complete recovery is 3-6 months.

4. General Instructions:

Apply ice to the shoulder (20min on /off) to reduce pain & swelling.

If you notice any signs of infection such as increased pain, redness, wound drainage or fever $>100.5^{\circ}$ F, call the office immediately.

5. Follow-up Appointment:

Call the office at (616) 949-8945 to schedule an appointment for 10-14 days after surgery.

6. Other Instructions:

*If you have any problems or concerns following surgery,
please call our office at (616) 949-8945.*